

(19)



Eur päisches Patentamt
Europ an Patent Office
Office européen d s brevets



(11) Publication number:

0 415 487 A2

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90202269.8

(51) Int. Cl.⁵: C07D 205/08, C07D 401/12,
C07D 405/06, A61K 31/395


(22) Date of filing: 24.08.90

(30) Priority: 31.08.89 US 401391

(43) Date of publication of application:
06.03.91 Bulletin 91/10(84) Designated Contracting States:
CH DE FR GB IT LI NL(71) Applicant: MERCK & CO. INC.
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065-0900(US)(72) Inventor: Yang, Shu Shu
324 Eileen Way
Bridgewater, NJ 08807(US)
Inventor: Chiang, Yuan-Ching P.
73 Mitchell Avenue
Piscataway, NJ 08854(US)
Inventor: Heck, James V.
961 Nepawin Lane
Scotch Plains, NJ 07076(US)
Inventor: Chang, Michael N.
2970 Windy Bush Road
Newton, PA 18940(US)(74) Representative: Hesketh, Alan, Dr. et al
European Patent Department Merck & Co.,
Inc. Terlings Park Eastwick Road
Harlow Essex, CM20 2QR(GB)

(54) Beta-lactams as anticholesterolemic agents.

(57) Compounds of formula (I), which are useful as antihypercholesterolemic agents, are disclosed.



(I)


EP 0 415 487 A2

β -LACTAMS AS ANTICHOLESTEROLEMIC AGENTS**BACKGROUND OF THE INVENTION**

Hypercholesterolemia is known to be one of the prime risk factors for ischemic cardiovascular disease, such as arteriosclerosis. Bile acid sequestrants have been used to treat this condition; but they seem to be moderately effective but they must be consumed in large quantities, i.e., several grams at a time, and they are not very palatable.

MEVACOR® (lovastatin), now commercially available is one of a group of very active antihypercholesterolemic agents that function by limiting cholesterol biosynthesis by inhibiting the enzyme, HMG-CoA reductase. Another approach to limiting cholesterol biosynthesis is through inhibition of the enzyme HMG-CoA synthase.

Copending U.S. patent application S.N. 053,774, filed May 26, 1987, corresponding to EP-A-0293132, discloses certain β -lactones of formula (i)




(1)

which are useful as antihypercholesterolemic agents and are believed to function by inhibiting HMG-CoA synthase. Additional β -lactones which have antihypercholesterolemic activity are disclosed in copending U.S. patent applications S.N. 021,848, filed March 4, 1987, corresponding to EP-A-0285254, and S. N. 053,646, filed May 26, 1987, corresponding to EP-A-0293128.

DETAILED DESCRIPTION OF THE INVENTION

The present invention is directed to compounds of structural formula (I):




(I)

wherein:

Q is

- a. C₁₋₅ alkyl;
- b. C₆₋₁₀ aryl or C₆₋₁₀ heteroaryl including one heteroatom selected from N, O, or S;
- c. C₇₋₁₅ aralkyl or C₇₋₁₅ heteroaralkyl;
- d. C₆₋₁₀ aryl or C₆₋₁₀ heteroaryl substituted with W;
- e. C₇₋₁₅ aryl or C₇₋₁₅ heteroaryl wherein the aryl or heteroaryl moiety is substituted with W;
- f. OH; R₂ and R₄ are independently selected from:


- a. H,
 b. C₁₋₅ hydroxyalkyl,
 c. C₁₋₅ alkyl,
 d. C₁₋₅ alkoxy,
 5 e. C₂₋₆ alk nyl, R₄ is CHR₀R;
 R₀ is H, C₁₋₅ alkyl, phenyl or phenyl C₁₋₅ alkyl;
 R is
 a. O-(CO)_{p2}-(A)_{p1}-R₁,
 b. OSO₂R₁,
 10 c. AR₁,
 d. R together with R₀ and the C to which it is attached form



- e. R together with R₀ and the C to which it is attached form




- p₁ and p₂ are independently 0 or 1; provided that p₁ and p₂ are not both 0;
 25 R₁ is
 a. H,
 b. aryl, or a heteroaryl group of 5 to 10 atoms, including one to two heteroatoms, selected from:



- c. phenyl S(O)_n, wherein n is 0 to 2,
 d. C₃₋₆ cycloalkyl; B is O, N or S;
 A is
 a.



b.



5

c. NH; R₂ is

a. H,

10

b. C₁₋₅ alkoxy,c. C₁₋₅ alkyl,

d. OH,

e. OC(O)CH₃; W isa. C₁₋₅ alkyl,

15

b. C₁₋₅ alkoxy,c. NO₂,d. NHR₃,e. C₁₋₃ alkylS(O)p₃

f. halogen,

20

g. COOR₃,h. OCOR₃,i. NHCOR₃,j. NHSO₂R₃,k. NHSO₂phenyl; X, Y, and Z are independently selected from:

25

a. H,

b. halogen,

c. C₁₋₅ alkyld. C₁₋₅ alkoxy,e. C₁₋₅ alkoxy carbonyl

30

f. C₁₋₅ alkyl carbonyloxyamino,

g. amino,

h. phenoxy,

i. phenyl C₁₋₅ alkoxy,j. C₁₋₅ alkylSO₂NH,

35

k. NO₂,

l. diphenylmethyloxycarbonyl,

m. C₁₋₅ alkylthio,

n. aminocarbonyl,

o. carboxy,

40

p. C₁₋₅ alkenyloxyq. C₁₋₅ alkylamido,

r. trifluoromethyl.

One embodiment of the present invention are those compounds of formula (I) wherein:
R is O-(CO)_{P2}-(A)_{P1}-R₁ or OSO₂R₁, and R₀ and R₅ are H.

45


In one class of this embodiment are those compounds wherein R is O-(CO)_{P2}-(A)_{P1}-R₁.

In one subclass are those compounds wherein P₁ is 0 and P₂ is 1 and Q is 4-methylphenyl or OH.
Exemplifying this subclass are the compounds of formula (I) described in Table I.

50

55

TABLE I



CONFIGURATION

AT 3, 4

POSITIONSR₃R₄IC₅₀

a.	<u>cis</u>	CH ₃ CH ₂ -	3-thienylCO ₂ CH ₂ -	7.1 x 10 ⁻⁷
b.	<u>cis</u>	CH ₃ CH ₂ -	3-methanesulfonamidophenyl-CO ₂ CH ₂ -	2.5 x 10 ⁻⁸
c.	<u>cis</u>	CH ₃ CH ₂ -	2-nitrophenylCO ₂ CH ₂ -	2.3 x 10 ⁻⁸
d.	<u>cis</u>	CH ₃ CH ₂ -	4-pyridylCO ₂ CH ₂ -	6.2 x 10 ⁻⁸
e.	<u>cis</u>	CH ₃ CH ₂ -	2-pyridylCO ₂ CH ₂ -	5.4 x 10 ⁻⁸
f.	<u>cis</u>	CH ₃ CH ₂ -	5-(2-methoxypyridyl)CO ₂ CH ₂ -	4.1 x 10 ⁻⁸
g.	<u>cis</u>	CH ₃ CH ₂ -	2-furanylCO ₂ CH ₂ -	7.9 x 10 ⁻⁹
h.	<u>cis</u>	CH ₃ CH ₂ -	2-(5-n-butylpyridyl)CO ₂ CH ₂ -	4.5 x 10 ⁻⁸
i.	3S,4S	CH ₃ CH ₂ -	4-methoxyphenylCO ₂ CH ₂ -	3.0 x 10 ⁻⁸
j.	3R,4R	CH ₃ CH ₂ -	4-methoxyphenylCO ₂ CH ₂ -	2.0 x 10 ⁻⁷
k.	<u>cis</u>	CH ₃ CH ₂ -	3-nitrophenylCO ₂ CH ₂ -	3.9 x 10 ⁻⁸
l.	<u>cis</u>	CH ₃ CH ₂ -	2-CH ₃ SO ₂ NHphenylCO ₂ CH ₂ -	3.1 x 10 ⁻⁸
m.	<u>cis</u>	CH ₃ CH ₂ -	4-pyridazinylCO ₂ CH ₂ -	8.2 x 10 ⁻⁸
n.	<u>cis</u>	CH ₃ CH ₂ -	2-NH ₂ phenylCO ₂ CH ₂ -	3.1 x 10 ⁻⁸
o.	<u>cis</u>	CH ₃ CH ₂ -	3-furanylCO ₂ CH ₂ -	3.2 x 10 ⁻⁷
p.	<u>cis</u>	CH ₃ CH ₂ -	2-quinolinylnylCO ₂ CH ₂ -	2.1 x 10 ⁻⁸
q.	<u>cis</u>	CH ₃ CH ₂ -	4-benzoxypyphenylCO ₂ CH ₂ -	4.0 x 10 ⁻⁸
r.	<u>cis</u>	CH ₃ CH ₂ -	4-quinolinylnylCO ₂ CH ₂ -	2.5 x 10 ⁻⁸
s.	<u>cis</u>	CH ₃ CH ₂ -	2-thienylCO ₂ CH ₂ -	3.8 x 10 ⁻⁸
t.	<u>cis</u>	CH ₃ CH ₂ -	3-(2-phenoxyipyridyl)CO ₂ CH ₂ -	3.3 x 10 ⁻⁸

TABLE I

5

CONFIGURATION

AT 3, 4

10

POSITIONS

R_3

R_4

IC_{50}

u. cis CH_3CH_2- 3- CH_3CONH phenyl CO_2CH_2- 7.2×10^{-8}

v. cis CH_3CH_2- 3- NH_2 phenyl CO_2CH_2- 4.5×10^{-8}

w. cis CH_3CH_2- 4-((Ph) $_2COOC$)phenyl CO_2CH_2- 6.2×10^{-8}

x. cis CH_3CH_2- 4- $HOOC$ phenyl CO_2CH_2- 3.5×10^{-7}

15 y. cis CH_3CH_2- 4- NO_2 phenyl CO_2CH_2- 2.8×10^{-8}

z. cis $CH_3CH_2CH_2-$ 4- CH_3O phenyl CO_2CH_2- 6.95×10^{-8}

aa. cis $HOCH_2-$ 4- CH_3O phenyl CO_2CH_2- 1.8×10^{-7}

20 bb. cis CH_3CH_2- 3-pyridyl CO_2CH_2- 6.4×10^{-8}

cc. cis CH_3CH_2- 4- CH_3O phenyl CO_2CH_2- 3.0×10^{-8}

dd. cis CH_3CH_2- phenyl CO_2CH_2- 1.1×10^{-7}

ee. cis CH_3CH_2- 3-Coumaryl CO_2CH_2- 5.0×10^{-8}

25 ff. cis CH_3CH_2- 2-benzofuranyl CO_2CH_2- 3.3×10^{-8}

gg. S,S CH_3CH_2- 2-furanyl CO_2CH_2- 1.1×10^{-8}

hh. cis CH_3CHCH_3- 4-methoxyphenyl CO_2CH_2- 1.3×10^{-7}

30 ii. cis CH_3CH_2- 4- CH_3O_2C phenyl CO_2CH_2- 8.5×10^{-9}

jj. cis CH_3CH_2- 4-n-butylcarbamoylphenyl CO_2CH_2- 2.0×10^{-8}

kk. cis CH_3CH_2- 4-aminophenyl CO_2CH_2- 4.7×10^{-8}

35 ll. cis CH_3CH_2- 3-methoxy-4-allyloxy-5-nitro-phenyl CO_2CH_2- 1.6×10^{-7}

mm. cis CH_3CH_2- 3-methoxy-4-allyloxy-5-(methyl-amido)phenyl CO_2CH_2- 3.0×10^{-7}

40 nn. cis CH_3CH_2- 4-methanesulfonamidophenyl CO_2CH_2- 1.6×10^{-8}

oo. cis CH_3CH_2- 5-(2,3-dimethoxypyridyl) CO_2CH_2- 3.8×10^{-8}

pp. cis CH_3CH_2- 6-(2,3-dimethoxypyridyl) CO_2CH_2- 8.9×10^{-8}

45 qq. cis CH_3CH_2- 4-(2-methoxythiazolyl) CO_2CH_2- 4.5×10^{-8}


rr. trans $CH_3CH_2CH_2-$ 4- CH_3O phenyl CO_2CH_2- 3.5×10^{-8}

ss. trans $HOCH_2-$ 4- CH_3O phenyl CO_2CH_2- 1.5×10^{-7}

50 tt. trans CH_3CH_2- 4- CH_3O phenyl CO_2CH_2- 1.7×10^{-7}

In a second subclass are those compounds of formula (I) wherein p₁ is 1 and p₂ is 1. Exemplifying this subclass are the compounds of formula (I) described in Table II.

TABLE II



CONFIGURATION

AT 3, 4

	POSITIONS	R ₃	R ₄	IC ₅₀
a.	<u>cis</u>	CH ₃ CH ₂ -	4-CH ₃ OphenylC=CCO ₂ CH ₂ -	9.5 x 10 ⁻⁹
b.	<u>cis</u>	CH ₃ CH ₂ -	4-CH ₃ OphenylCH ₂ CO ₂ CH ₂ -	3.5 x 10 ⁻⁸
c.	<u>cis</u>	CH ₃ CH ₂ -	3-pyridylCH ₂ CO ₂ CH ₂ -	3.5 x 10 ⁻⁸
d.	3S,4S	CH ₃ CH ₂ -	phenyl-(R)-CHOCH ₃ CO ₂ CH ₂ -	1.8 x 10 ⁻⁸
e.	4S	H	phenyl-(S)-CHOCH ₃ CO ₂ CH ₂ -	4.2 x 10 ⁻⁸
f.	4S	H	phenyl-(R)-CHOCH ₃ CO ₂ CH ₂ -	5.0 x 10 ⁻⁸
g.	<u>cis</u>	CH ₃ CH ₂ -	phenyl-(R)-CHOCH ₃ CO ₂ CH ₂ -	6.7 x 10 ⁻⁸
h.	<u>cis</u>	CH ₃ CH ₂ -	3-pyridylC=CCO ₂ CH ₂ -	3.1 x 10 ⁻⁸
i.	3S,4S	CH ₃ CH ₂ -	3-pyridylCH ₂ CO ₂ CH ₂ -	8.7 x 10 ⁻⁹
j.	3S,4S	CH ₃ CH ₂ -	4-methoxyphenylCH ₂ CO ₂ CH ₂ -	8.6 x 10 ⁻⁹
k.	3S,4S	CH ₃ CH ₂ -	4-methoxyphenylC=CCO ₂ CH ₂ -	1.1 x 10 ⁻⁸
l.	<u>cis</u>	CH ₃ CH ₂ -	2-furanylCH ₂ CO ₂ CH ₂ -	2.4 x 10 ⁻⁶
m.	<u>cis</u>	CH ₃ CH ₂ -	2-furanylC=CCO ₂ CH ₂ -	2.5 x 10 ⁻⁷
n.	<u>cis</u>	CH ₃ CH ₂ -	phenoxyCH ₂ CO ₂ CH ₂ -	2.4 x 10 ⁻⁸

TABLE II

CONFIGURATION				
AT 3, 4				
	<u>POSITIONS</u>	<u>R₃</u>	<u>R₄</u>	<u>IC₅₀</u>
5	o. <u>sia</u>	CH ₃ CH ₂ -	3- acetyl -4- ethoxy phenylCH ₂ CO ₂ CH ₂ -	2.1 x 10 ⁻⁹
	p. <u>sia</u>	CH ₃ CH ₂ -	2-pyridylCH ₂ CO ₂ CH ₂ -	1 x 10 ⁻⁸
10	q. <u>sia</u>	CH ₃ CH ₂ -	phenylthioCH ₂ CO ₂ CH ₂ -	4.2 x 10 ⁻⁸
	r. <u>sia</u>	CH ₃ CH ₂ -	4- ethoxy phenylCH ₂ CO ₂ CH ₂ -	2.2 x 10 ⁻⁸
	s. <u>sia</u>	CH ₃ CH ₂ -	3,4,5-tri ethoxy phenylCH ₂ CO ₂ CH ₂ -	2.0 x 10 ⁻⁸
15	t. <u>sia</u>	CH ₃ CH ₂ -	3,4,5-tri ethoxy phenylC=CCO ₂ CH ₂ -	3.8 x 10 ⁻⁸
	u. <u>sia</u>	CH ₃ CH ₂ -	3-bromo-4-propanoxy-5- ethoxy - phenylC=CCO ₂ CH ₂ -	9 x 10 ⁻⁸
	v. <u>sia</u>	CH ₃ CH ₂ -	4- acetyl phenylaminoCO ₂ CH ₂ -	5.2 x 10 ⁻⁸
20	w. <u>sia</u>	CH ₃ CH ₂ -	cyclohexylaminoCO ₂ CH ₂ -	5.4 x 10 ⁻⁸
	x. <u>sia</u>	CH ₃ CH ₂ -	4-trifluoro acetyl phenylamino- CO ₂ CH ₂ -	2.6 x 10 ⁻⁷
25	y. <u>sia</u>	CH ₃ CH ₂ -	3,4-di ethoxy phenylCH ₂ CO ₂ CH ₂ -	3 x 10 ⁻⁸
	z. <u>sia</u>	CH ₃ CH ₂ -	6-(2,3-di ethoxy pyridyl)C=CCO ₂ CH ₂ -	2.7 x 10 ⁻⁸
	aa. <u>sia</u>	CH ₃ CH ₂ -	4-fluorophenylaminoCO ₂ CH ₂ -	5.0 x 10 ⁻⁸

30

Double bonds within the R₄ moiety are in the trans configurations.

In a second class of this embodiment are those compounds wherein R is OSO₂R₁. Exemplifying this
35 class is the compound described in Table III.

40

45

50


55

TABLE III

5

10

15



20

CONFIGURATION

AT 3, 4

POSITIONS R_3 R_4 IC_{50}


25

sis $CH_3CH_2-4-Cl_3phenylSO_2OCH_2-$ 1.1×10^{-7}

30

In a second embodiment of the present invention are those compounds of formula (I) wherein:
R is $O-(CO)_{P_2}-(A)_{P_1}-R_1$, AR_1 or R together with R_0 and the C to which it is attached form

35



P_1 and P_2 are independently 0 or 1, provided that P_1 and P_2 are not both zero;
 R_0 is H, CH_3 , or phenyl; provided that when R is $O-(CO)_{P_2}-(A)_{P_1}-R_1$, R_0 is CH_3 or phenyl;
 R_5 is H.

40

In one class of this embodiment are the compounds wherein:

45




R_0 is H, or CH_3 . Exemplifying this class are the compounds in Table IV.

50

55

TABLE IV



CONFIGURATION

AT 3,4

	<u>POSITIONS</u>	R_3	R_4	IC_{50}
a.	<u>trans</u>	$CH_3CH_2CH_2-$	$phenylC\equiv C-$	1.5×10^{-6}
b.	<u>cis</u>	$CH_3CH_2CH_2-$	$phenylC\equiv C-$	2.2×10^{-6}
c.	<u>trans</u>	$HOCH_2-$	$phenylC\equiv C-$	1.1×10^{-6}
d.	<u>cis</u>	$HOCH_2-$	$phenylC\equiv C-$	1.1×10^{-6}
e.	<u>trans</u>	CH_3CH_2-	$phenylC\equiv CCH_3$	5.0×10^{-5}
f.	<u>cis</u>	CH_3CH_2-	$phenylC\equiv CCH_3$	4.3×10^{-7}


In a second class are the compounds wherein:

R is $O-(CO)_{P_2}-(A)_{P_1}-R_1$ or AR_1 ;

R_0 is H, CH_3 or phenyl; provided that when R is $O-(CO)_{P_2}-(A)_{P_1}-R_1$, R_0 is CH_3 or phenyl.

Further illustrating this class are those compounds wherein P_1 is O. Exemplifying this class are the compounds of table V.

TABLE V



CONFIGURATION

AT 3, 4



	<u>POSITIONS</u>	<u>R₃</u>	<u>R₄</u>	<u>IC₅₀</u>
a.	<u>cis</u>	CH ₃ CH ₂ -	4-HO ₂ CphenylNHCH ₂ -	9.9 x 10 ⁻⁶
b.	<u>cis</u>	CH ₃ CH ₂ -	phenylCH ₂ CH ₂ -	8.6 x 10 ⁻⁸
c.	<u>cis</u>	CH ₃ CH ₂ -	4-CH ₃ OphenylCO ₂ CHCH ₃	1.2 x 10 ⁻⁷
d.	<u>cis</u>	CH ₃ CH ₂ -	phenylCH ₂ CHCH ₃	1.75 x 10 ⁻⁷
e.	<u>cis</u>	CH ₃ CH ₂ -	4-CH ₃ OphenylCO ₂ CHphenyl	2 x 10 ⁻⁶
f.		H	phenylCHCHCH ₃ OCOCH ₃	4.9 x 10 ⁻⁷
g.		H	CH ₃ CO ₂ CH ₂ CHCH ₃	2.9 x 10 ⁻⁷

A third embodiment of the present invention is represented by those compounds of formula (I) wherein:

R₀ is H, CH₃ or CH₂phenyl;


R₅ is C₁₋₅ alkyl, or C₁₋₅ alkoxy; and


R is O-(CO)p₂-(A)p₁-R₁, AR₁, or R together with R₀ and the C to which it is attached form



P₁ and P₂ are independently 0 or 1; provided that both P₁ and P₂ are not both zero. Exemplifying this embodiment are those compounds of formula (I) described in Table VI.

TABLE VI



	R ₃	R ₅	R ₄	IC ₅₀
20	a. CH ₃	CH ₃	phenylCH ₂ CH ₂ -	1.8 x 10 ⁻⁶
	b. CH ₃	CH ₃	phenylCO ₂ CHCH ₂ phenyl	4 x 10 ⁻⁷
	c. CH ₃	CH ₃	phenylCH=CH-	6.8 x 10 ⁻⁷
25	d. CH ₃	CH ₃	phenylCH-CH- 	1.1 x 10 ⁻⁷

The alkyl groups referred to above may be straight chain or branched or may include cycloalkyl groups.

Halogen or halo means fluoro, chloro, bromo or iodo.

The non-optically active compounds of the present invention may be prepared according to Scheme I. Trimethylsilyl amine is added to an α,β unsaturated aldehyde to form an imine which is then condensed with an ester moiety for form a β -lactam of formula (I-1). Compound (I-1) may, under ozonolysis conditions, be converted to the aldehyde (I-2) or the alcohol (I-3) or (I-1) may be reduced to compound (I-4) which can be sulfonated at the lactam nitrogen. Compound (I-1) may also be converted to the epoxide which can be converted to esters of formula (I-6). The aldehyde compound (I-2) may be treated with a Grignard reagent to form compounds of formula (I-7) where R₆ is methyl or phenyl or benzyl. The alcohol (I-3) can be protected at the hydroxyl group, then sulfonated, deprotected and treated with an acylating agent to form compounds of formula (I-9) or an isocyanate to form compounds of formula (I-10). In the alternative the acylation at the hydroxyl group may precede the sulfonation at the lactam nitrogen.

Substituents on an aryl moiety either in R' or R'' can be formed after acylation or sulfonation by conversion of aryl substituents such as -NO₂ to -NH₂ followed by further conversion of amino to an amide or sulfonamide functionality.

SCHEME I

5

10

15

20

25

30


35


40

45

50

55





The optically active compounds of the present invention can be prepared as outlined in Scheme II.

SCHEME II

5

10

15

20

25

30


35

40

45

50

55



Where the *cis* R,R stereoisomer is desired it may be formed in the above sequence by employing D-aspartic acid in place of L-aspartic acid. The alcohol (II-1) may be further converted to the corresponding alkyl halide with retention of configuration.

The present invention is also directed to a method of inhibiting cholesterol biosynthesis which comprises the administration to a subject in need of such treatment a nontoxic, therapeutically effective amount of a compound represented by the following general structural formula (I) and pharmaceutically acceptable salts thereof.

The present invention is also directed to a method of inhibiting the activity of HMG-CoA synthase enzyme which comprises the administration to a subject in need of such treatment a nontoxic, therapeutically effective amount of a compound represented by the general structural formula (I) and pharmaceutically acceptable salts thereof.

Specifically the compounds of this invention are useful as antihypercholesterolemic agents for the treatment of arteriosclerosis, hyperlipidemia, familiar hypercholesterolemia and the like diseases in humans. They may be administered parenterally in the form of a capsule, a tablet, an injectable preparation or the like. Doses may be varied, depending on the age, severity, body weight and other conditions of human patients but daily dosage for adults is within a range of from about 20 mg to 2000 mg (preferably 20 to 100 mg) which may be given in two to four divided doses. High doses may be favorably employed as required.

The pharmaceutically acceptable salts of the compounds of this invention include those formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc, and from bases such as ammonia, ethylenediamine, N-methylglucamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chloroprocaine, diethanolamine, procaine, N-benzylphenethylamine, diethylamine, piperazine, tris(hydroxymethyl)aminomethane, and tetramethylammonium hydroxide.

The compounds of this invention may also be coadministered with pharmaceutically acceptable nontoxic cationic polymers capable of binding bile acids in a non-reabsorbable form in the gastrointestinal tract. Examples of such polymers include cholestyramine, colestipol and poly[methyl-(3-trimethylaminopropyl)imino-trimethylene dihalide]. The relative amounts of the compounds of this invention and these polymers is between 1:100 and 1:15,000.

The intrinsic HMG-CoA synthase inhibition activity of the compounds of this invention is measured by the standard *in vitro* protocol described below:

The livers from male Charles River CD rats (225-350 g) were homogenized in 0.25 M sucrose which was adjusted with phenylmethylsulfonyl fluoride (PMSF) and N-p-tosyl-L-lysine chloromethyl ketone (TLCK) so that the final concentration of each was 50 and 25 mg/ml, respectively. The homogenate was centrifuged at 15,000 x g for 20 minutes, the supernatant filtered through a fine nylon screen to remove most of the fat layer and recentrifuged at 100,000 x g for 1 hour. This supernatant was removed and 1 M potassium phosphate, dithiothreitol (DTT) and ethylene glycolbis(β -aminoethyl ether-N,N,N',N'-tetraacetic acid (EGTA) added to give a final concentration of 0.1 M (pH 7.2), 0.5 mM and 0.1 mM, respectively. Solid ammonium sulfate was added to 50% saturation to the protein solution, it was centrifuged at 15,000 x g and the supernatant discarded. This precipitated protein could be stored at -70 °C for at least one month with very little loss of activity. The ammonium sulfate precipitate was dissolved in an minimal amount of 0.06 M potassium phosphate buffer (pH 7.2) containing 0.5 mM dithiothreitol and 0.1 mM EGTA (referred to as 0.06 M phosphate buffer) and dialyzed overnight against 2 liters of the same buffer to remove the ammonium sulfate and to inactivate HMG-CoA lyase [Clinkenbeard, et al., J. Biol. Chem. 250, 3108-3116(1975)].

The dialyzed extract was added to a column of DEAE-52 (Whatman) which had been equilibrated with 0.06 M phosphate buffer (10 mg of protein to 1 ml bed volume of the resin). The DEAE-cellulose was eluted with 0.06 M phosphate buffer until the optical density at 280 nm was essentially zero. This fraction contained the β -ketoacetyl-CoA thiolase activity. The HMG-CoA synthase was eluted from the column with 0.1 M phosphate buffer (pH 7.2) containing 0.5 mM DTT and 0.1 mM EGTA, and was virtually free of all thiolase activity. The protein was precipitated by the addition of ammonium sulfate to give 50% saturation. This solution was stirred for 10 minutes at 4 °C and the precipitate collected by centrifugation at 15,000 rpm for 10 minutes. The supernatant was discarded and the precipitate dissolved in a minimum of 0.06 M phosphate buffer, pH 7.2 (about 10 ml) and the enzyme stored at -80 °C.

HMG-CoA Synthase Inhibition Assay

Enzyme protein (ca. 24 mg) was added to a solution containing 117 μ M Tris-HCl (pH 8.0), 11.7 μ M MgCl₂, 1.17 μ M ethylenediaminetetraacetic acid (EDTA), 0.58 μ M dithiothreitol, and the indicated concentrations of the test compound (added as a 2 mg/ml solution in dimethylsulfoxide). The incubation took place in a volume of 0.085 ml at 30 ° in a shaking water bath. After 5 minutes, 15 ml of a solution containing acetoacetyl-CoA and 0.1 μ Ci of 1-[¹⁴C]-acetyl-CoA was added to give a final concentrations of 0.1 and 0.4 μ M, respectively. The incubation was continued for 10 more minutes and the reaction stopped by the addition of 50 ml of the assay mixture to 0.2 ml of 6N HCl in a glass scintillation vial. The vial was heated for 1 hour at 120 ° after which time 0.2 ml more of 6N HCl was again added to each vial and the heating continued for another hour. Following this, 1.0 ml of 0.9% saline was added to each vial and finally 10 ml of scintillation liquid. Radioactivity was determined in a Packard Tri-Carb liquid scintillation counter. Percent inhibition is calculated by the formula:

$$1 - \frac{\text{Sample} - \text{Blank}}{\text{Control} - \text{Blank}}$$

IC₅₀ values were determined by plotting the log of the concentration of the test compound verses the percentage inhibition and fitting a straight line to the resulting data by using the least squares method.

The following examples illustrate the preparation of compounds of formula (I) and as such are not to be

considered as limiting the invention set forth in the claims appended hereto.

Example 1

Preparation of (±)-cis-3-Ethyl-4-hydroxymethyl-2-azetidinone

Method I:

Step A: Generation of N-Trimethylsilyl Imine Solution:

To a cooled solution of 10.54 ml (50 mmol) of 1,1,1,3,3,3- hexamethyldisilazane in 36 ml of anhydrous THF at -78°C was added 19.24 ml (48.1 mmol) of 2.5 M n -butyllithium in hexane in one portion. The resulting solution was stirred for 30 minutes at -78°C and to this solution was then added dropwise 5.7 ml (45.4 mmol) of cinnamaldehyde in 12 ml of THF. The mixture was stirred for 1 1/2 hours and the resulting cold solution of N-trimethylsilyl imine was used directly in the following reaction.

Step B: Preparation of (±)-cis-3-Ethyl-4-(β-styryl)-2-azetidinone

To a cooled solution of 7.26 ml (51.7 mmol) of diisopropylamine in 40 ml of THF at -78°C was added 21 ml (52.5 mmol) of 2.5 M n -butyllithium. The solution was stirred for 10 minutes at -78°C then 15 minutes at room temperature followed by the addition of 6 ml (45.4 mmol) of ethyl butyrate in 12 ml of THF. After stirring for 1 hour at -78°C , to this solution was added dropwise the above generated (Step 3A) cooled solution of N-trimethylsilyl imine. The resulting mixture was stirred at -78°C for 1 hour then at room temperature for 2 hours, quenched with 300 ml of cooled 1 N HCl, and extracted with 3 x 100 ml ether. The ether phases were combined, dried and concentrated. The product was purified by flash column chromatography (R_f = 0.25, 30% EtOAc in hexane) to afford the product β-lactam. NMR (CDCl_3): δ 1.00 (3H, t), 1.48-1.88 (2H, m), 3.32 (1H, m), 4.40 (1H, m), 6.12 (1H, s), 6.24 (1H, dd), 6.64 (1H, d), 7.21-7.38 (5H, m).

Step C: Preparation of (±)-cis-3-Ethyl-4-hydroxymethyl-2-azetidinone

A solution of 13 g (64.5 mmol) of (±)-cis -3-ethyl-4-(β-styryl)-2-azetidinone in 400 ml of CH_2Cl_2 was ozonized at -78°C until the solution turned blue. The resulting solution was stirred for 20 50 minutes at -78°C then 30 minutes at room temperature. The solution was recooled to -78°C and to this solution was added 14 ml (190 mmol) of methyl sulfide. After stirring for 10 minutes at -78°C and minutes at room temperature, the solution was concentrated and the residue was redissolved in 400 ml of methanol and cooled to 0°C . 5.5 g (149 mmol) of sodium borohydride was added slowly to the cooled solution. The resulting mixture was stirred for 20 minutes at 0°C then 1/2 hour at room temperature and concentrated. The product was purified by flash column chromatography to yield the titled compound. NMR (CDCl_3): δ 1.08 (3H, t), 1.50-1.89 (2H, m), 3.17 (1H, m) 3.66-4.00 (3H, m), 6.89 (1H, s).

Method II:

To a solution of 500 mg of (±)-cis -3-ethyl-4-benzyloxycarbonyl-2-azetidinone (2.1 mmol) in 20 ml of anhydrous THF was added dropwise 1.0 ml of 2M lithium borohydride-THF solution. The mixture was stirred at room temperature for 5 hours and quenched with 1.5 ml of methanol dropwise in the cold. The mixture was concentrated and the residue purified by preparative tlc on silica gel plates developed with ethyl acetate to give the titled compound.

Example 25 Preparation of (±)-cis-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinoneStep A: Preparation of (±)-cis-3-Ethyl-4-t-butyldimethylsilyloxymethyl-2-azetidinone

10 187 mg of (±)-cis-3-ethyl-4-hydroxymethyl-2-azetidinone, 300 mg of imidazole, and 300 mg of t-butyldimethylsilyl chloride in 3 ml dry dimethylformamide (DMF) were stirred at room temperature for 2 hours. The mixture was concentrated to dryness and 25 ml of H₂O:EtOAc (2:3) were added. The mixture was washed 3 times with 3 ml of H₂O to remove DMF and imidazole and then dried over Na₂SO₄ and concentrated to yield the titled compound as white crystals. NMR (CDCl₃): δ 0.06 (6H, s), 0.89 (9H, s), 1.06 (3H, t), 1.48-1.88 (2H, m), 3.14 (1H, m), 3.56-3.90 (3H, m), 5.86 (1H, brs).

Step B: Preparation of (±)-cis-3-Ethyl-4-t-butyldimethylsilyloxymethyl-N-toluenesulfonyl-2-azetidinone

20 The product of Step 2A and 300 mg of tosyl chloride were dissolved in 10 ml of 1% tetrabutylammonium bromide solution (CH₃CN:CH₂Cl₂ = 1:19). 100 mg KOH was added and the mixture stirred for one hour. The mixture was filtered and the filtrate concentrated to give the crude product. NMR (CDCl₃): δ 0.07 (6H, d), 0.88 (9H, s), 1.04 (3H, t), 1.8 (2H, m), 2.44 (3H, s), 3.15 (1H, m), 3.84-4.16 (3H, m), 7.4 (2H, d), 7.9 (2H, d).

25

Step C: Preparation of (±)-cis-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone

30 To the product of Step 2B (100 mg) dissolved in 2 ml of THF was added 150 µl of aqueous HF, and the mixture stirred at room temperature for 4 hours. The mixture was filtered through silica and washed with 1:1 EtOAc/aq. NaHCO₃ and dried over Na₂SO₄. The mixture was filtered and concentrated to a colorless oil which was flash chromatographed to yield the title compound. NMR (CDCl₃): δ 1.04 (3H, t), 1.50-1.92 (2H, m), 2.47 (3H, s), 2.65 (1H, brs), 3.15 (1H, dt), 3.80-4.15 (3H, m), 7.40 (2H, d), 7.90 (2H, d).

35

Example 340 Preparation of (±)-cis-3-Ethyl-4-p-methoxyphenylacetoxymethyl-N-toluenesulfonyl-2-azetidinone

To (±)-cis-3-ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone (3.3 mg) and p-methoxyphenylacetic acid (3.7 mg) in CH₂Cl₂ (0.6 ml) was added dicyclohexylcarbodiimide (DCC) (5 mg) followed by p-dimethylaminopyridine (DMAP) (1 mg). The mixture was stirred overnight, at room temperature and then purified via preparative tlc on a silica gel plate and developed with CH₂Cl₂ to yield the titled compound. NMR (CDCl₃): δ 0.94 (3H, t), 1.22-1.74 (2H, m), 2.44 (3H, s), 3.18 (1H, dt), 3.45 (2H, q), 3.79 (3H, s), 4.20-4.50 (3H, m), 6.98 (4H, dd), 7.61 (4H, dd).

50

Example 4Preparation of (±)-cis-3-Ethyl-4-(2-phenyl)ethyl-2-azetidinone

55

- To (±)-cis-3-ethyl-4-(β-styryl)-2-azetidinone (28 mg) in 2 ml of EtOAc was added 3 mg of 10% Pd/C. The mixture was hydrogenated at room temperature and 1 atmosphere. (H₂ consumed = 0.000139 moles). The solution was filtered and the filtrate concentrated and purified by preparative tlc, to yield the titled

compound. NMR (CDCl₃): δ 1.06 (3H, t), 1.45-2.08 (4H, m), 2.70 (2H, dt), 3.11 (1H, dt), 3.70 (1H, m), 5.70 (1H, brs), 7.10-7.39 (5H, m).

Example 5

Preparation of cis-(3S),(4S)-3-Ethyl-4-[(R)- α -methoxyphenylacetoxy]methyl-N-toluenesulfonyl-2-azetidinone
 and cis-(3R),(4R)-3-Ethyl-4-[(R)- α -methoxyphenylacetoxy]methyl-N-toluenesulfonyl-2-azetidinone

Step A: Preparation of cis(\pm)-3-Ethyl-4-[(R)- α -methoxyphenylacetoxy]methyl-2-azetidinone

To (\pm)-cis -3-ethyl-hydroxymethyl-2-azetidinone (26 mg) and R-(-)- α -methoxyphenylacetic acid (54 mg) in CH₂Cl₂ (6 ml) was added DCC (62 mg) followed by addition of p -dimethylaminopyridine (12 mg). The mixture was stirred at room temperature overnight and the filtrate concentrated and purified via preparative tlc on a silica gel plate (1500 μ) and developed with CH₂Cl₂ halfway and then EtOAc:CH₂Cl₂ = 1:1 to yield the titled compound. NMR (CDCl₃): δ 1.00 and 1.01 (3H, 2t), 1.2-1.8 (2H, m), 3.15 (1H, m), 3.21 (3H, s), 3.79 (1H, m), 4.15 (1H, m), 4.38 (1H, m), 4.80 (1H, s), 5.61 (1H, brs), 7.40 (5H, m).

Step B: Preparation of cis(\pm)-3-Ethyl-4-[(R)- α -methoxyphenylacetoxy]methyl-N-toluenesulfonyl-2-azetidinone

To the product of Step 5A (20 mg) in 0.8 ml of 1% tetrabutyl ammonium bromid solution (CH₃CN:CH₂Cl₂, 1:19) (TBAB solution) was added excess tosyl chloride and pulverized KOH. The mixture was stirred at room temperature for 36 hours. The mixture was then filtered and concentrated and the residue purified by preparative tlc on a silica gel plate (1500 μ) developed halfway with CH₂Cl₂ and then 5-10% EtOAc in CH₂Cl₂, to yield the titled compound. NMR (CDCl₃): δ 0.78 and 0.88 (3H, 2t), 1.00-1.70 (2H, m), 2.45 and 2.47 (3H, 2s), 3.1 (1H, m), 3.38 and 3.42 (3H, 2s), 4.10-4.38 (2H, m), 4.40-4.60 (1H, m), 4.49 and 4.80 (1H, 2s), 7.20-7.97 (9H, m).

Step C: Separation of cis-(3S),(4S)-3-Ethyl-4-[(R)- α -methoxyphenylacetoxy]methyl-N-toluenesulfonyl-2-azetidinone and cis-(3R),(4R)-3-Ethyl-4-[(R)- α -methoxyphenylacetoxy]methyl-N-toluenesulfonyl-2-azetidinone

The product of Step 5B (3.6 mg) in CH₂Cl₂ (0.25 ml) was spotted dropwise on 2 silica gel plates (250 μ) and developed with:
 EtOAc:Hexane = 1:9 4 times
 EtOAc:Hexane = 1:8 2 times
 EtOAc:Hexane = 1:7.5 once
 to give the (3S),(4S) titled compound (high R_f); NMR (CDCl₃): δ 0.88 (3H, t), 1.14-1.70 (2H, m), 2.45 (3H, s), 3.1 (1H, m), 3.42 (3H, s), 4.12-4.38 (2H, m), 4.48-4.60 (1H, m), 4.80 (1H, s), 7.20-7.90 (9H, m); and the (3R),(4R) titled compound (low R_f); NMR (CDCl₃): δ 0.78 (3H, t), 1.00-1.78 (2H, m), 2.47 (3H, s), 3.1 (1H, m), 3.38 (3H, s), 4.10-4.38 (2H, m), 4.40-4.60 (1H, m), 4.49 (1H, s), 7.20-7.96 (9H, m).

Example 6

Preparation of cis-(3S),(4S)-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone and trans-(3R),(4S)-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone

Step A: Preparation of (3S)-3-Benzoyloxycarbonylamino-2-ethyl- γ -butyrolactone

The titled compound was prepared from N-CBZ-L-aspartic acid in three steps via the formation of N-

CBZ-L-aspartic acid anhydride and (3S)-3-benzyloxycarbonylamino- γ -butyrolactone according to the published procedure by Y. Takahashi et al., Chem. Pharm. Bull., 34, 3020 (1986).

5 Step B: Preparation of (3S)-3-Benzyloxycarbonylamino-4-t-butyldimethylsilyloxy-2-ethylbutyric acid

To a solution of (3S)-3-Benzyloxycarbonylamino-2-ethyl- γ -butyrolactone (24 g) in 300 ml of methanol was slowly added 100 ml of 1 N NaOH at 0 °C. The mixture was stirred at 0-5 °C for 25 minutes and then at room temperature for 20 minutes. Methylene chloride was added (2 x 150 ml) followed by the addition of
10 2.5 N HCl to bring the pH of the aqueous layer to 6.0. Ethyl acetate (900 ml) was added to the aqueous solution and the EtOAc layer concentrated to an off-yellow oil. The aqueous layer was brought to pH = 3.0 and then extracted again with EtOAc. The two EtOAc extracts were combined and dried over Na₂SO₄ and concentrated to a yellow oil which then crystallized in an ice bath.

The crystallized product was combined with 75 g t-butyldimethylsilyl chloride and 70 ml of triethylamine and 400 ml of DMF and stirred for 2 hours. The precipitated salts were filtered and the filtrate
15 stirred for 30 minutes and then concentrated to 100 ml. An ethyl ether:water (1000 ml:200 ml) mixture was added and the ether layer concentrated to a yellow oil which was then dissolved in 500 ml of methanol. The mixture was stirred at 0 °C and 100 ml of 1 N HCl added and stirred for 15 minutes at 0 °C and 10 minutes at room temperature. 80 ml of 1 N NaOH was added to bring the pH to 5.0 and the mixture concentrated to
20 200 ml. The EtOAc layer was washed with sat. NaCl and dried and concentrated to a brown oil. The crude solid was purified by flash chromatography to yield the titled compound. NMR (CDCl₃): δ 0.07 (6H, s), 0.86 (9H, s), 0.80-1.12 (2H, m), 1.5-1.9 (2H, m), 2.56 (1H, m), 3.68 (1H, m), 3.98 (1H, m), 4.25-4.50 (1H, m), 5.12 (2H, s), 5.26 (1H, d), 7.36 (5H, s).

25

Step C: Preparation of (3S)-3-Amino-4-t-butyldimethylsilyloxy-2-ethylbutyric Acid

To a solution of the product of Step 6B in 500 ml of methanol was added 2.0 grams of 10% Pd/C and the mixture hydrogenated over hydrogen at 40 psi. The catalyst was removed by filtration and the solution
30 concentrated to a colorless oil. The oil crystallized in ether/hexane. The solid was broken up, filtered and washed with ether to yield the titled compound which was employed in the next step.

Step D: Preparation of (4S)-4-t-Butyldimethylsilyloxymethyl-3-ethyl-2-azetidinone

35

The product from Step 6C (8.0 g), 11.0 g of triphenylphosphine, 8.8 g of 2,2'-dipyridyl disulfide in 3 liters of CH₃CN were heated for 4 hours. The mixture was concentrated to dryness, redissolved in CH₂Cl₂ and stirred at 0 °C with the addition of 16 ml of Et₃N and 20 ml of CH₃I. The mixture was stirred for 30 minutes at 0 °C and concentrated to dryness. Ether (500 ml) was added and the insoluble salts filtered off.
40 The ether solution was concentrated to a reddish oil and the crude product flash chromatographed using the following solvent gradient:

1 l 30% CH₂Cl₂/hexane

1 l 50% CH₂Cl₂/hexane

1 l CH₂Cl₂

45 500 ml ether

to yield the titled product.

NMR (CDCl₃): δ 0.06 (6H, s), 0.88 (9H, s), 1.06 (3H, t), 1.50-1.86 (2H, m), 3.14 (1H, m), 3.58-3.86 (3H, m), 5.86 (1H, brs).

50

Step E: Preparation of (4S)-4-t-Butyldimethylsilyloxymethyl-3-ethyl-N-toluenesulfonyl-2-azetidinone

The product of Step 6D (2.0 g), 20 g of tosyl chloride and 1.2 g of powdered KOH were stirred in 120 ml of 5% CH₃CN in CH₂Cl₂ containing 1% tetrabutylammonium bromide, at 0 °C for 15 minutes, then room
55 temperature for 90 minutes. The mixture was chromatographed (CH₂Cl₂:hexane:ether = 3:7:1) to yield a 3:1 cis : trans mixture. NMR (CDCl₃): δ 0.07 (6H, s), 0.87 (9H, s), 1.05 (3H, t), 1.58-1.94 (2H, m), 2.45 (3H, s), 3.15 (1H, m), 3.82-4.14 (3H, m), 7.36 (2H, d), 7.88 (2H, d).

Step F: Preparation of cis-(3S),(4S)-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone and trans-(3R),(4S)-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone

The mixture from Step 8E was dissolved in 50 ml of THF and 1.0 g of tetrabutylammonium fluoride and 2.0 ml of 50% HF were added. The mixture was stirred at 0 °C for 30 minutes, then room temperature for 18 hours. Saturated NaHCO₃ solution was added to neutralize the reaction mixture followed by ether/H₂O with stirring for 30 minutes. The mixture was filtered, concentrated to dryness and dissolved in ether/EtOAc. The mixture was flash chromatographed to yield the cis (S,S) and trans (R,S) isomers.
 NMR of cis isomer (CDCl₃): δ 1.04 (3H, t), 1.5-1.9 (2H, m), 2.47 (3H, s), 2.70 (1H, dd), 3.15 (1H, dt), 3.82-4.12 (3H, m), 7.38 (2H, d), 7.90 (2H, d).
 NMR of trans isomer (CDCl₃): δ 0.92 (3H, t), 1.32-1.94 (2H, m), 2.46 (3H, s), 2.49 (1H, m), 3.80-4.30 (3H, m), 4.78 (1H, m), 7.36 (2H, d), 7.78 (2H, d).

Example 7

Preparation of cis-(3S),(4S)-3-Ethyl-4-m-pyridine acetoxymethyl-N-toluenesulfonyl-2-azetidinone

cis -(3S),(4S)-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone (137 mg) and 70 mg of 3-pyridylacetic acid, 125 mg of DCC, and 5 mg of DMAP in 5 ml CH₂Cl₂ were stirred at room temperature for 18 hours. The mixture was worked up and the product crystallized from EtOAc/hexane.
 NMR (CDCl₃): δ 0.97 (3H, t), 1.3-1.8 (2H, m), 2.44 (3H, s), 3.20 (1H, dt), 3.60 (2H, s), 4.2-4.6 (3H, m), 7.2-7.9 (8H, m).

Example 8

Preparation of cis-(3R),(4R)-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone and trans-(3S),(4R)-3-Ethyl-4-hydroxymethyl-N-toluenesulfonyl-2-azetidinone

Using N-CBZ-D-aspartic acid in place of N-CBZ-L-aspartic acid and employing the procedure of Example 6 the titled compounds were obtained.

Example 9

Preparation of (±) cis-3-Ethyl-4-hydroxycarbonylphenylaminomethyl-N-toluenesulfonyl-2-azetidinone

Step A: Preparation of (±)-cis-3-Ethyl-4-(β-styryl)-N-toluenesulfonyl-2-azetidinone

70 mg of (±)-cis -3-Ethyl-4-(β-styryl)-2-azetidinone was dissolved in 3 ml of TBAB solution; 100 mg of tosyl chloride and 20 mg KOH were added and stirred at 0 °C for 15 minutes, and then room temperature for one hour. The crude product was chromatographed on a silica column using 20% EtOAc in hexanes as eluent to give the titled compound.
 NMR (CDCl₃): δ 0.96 (3H, t), 1.40-1.86 (2H, m), 2.43 (3H, s), 3.32 (1H, m), 4.76 (1H, m), 5.92 (1H, dd), 6.73 (1H, d), 7.29 (7H, m), 7.81 (2H, d).

Step B: Preparation of (±)-cis-3-Ethyl-4-formyl-N-toluenesulfonyl-2-azetidinone

The product of step (9A) was dissolved in 5 ml of a 2:1 mixture of CH₂Cl₂ and methanol. The mixture

was stirred at -78°C and ozone was bubbled in until a blue color persisted. The mixture was stirred at -78°C for 30 minutes and then room temperature for 30 minutes. Dimethyl sulfide (200 μl) was added and the mixture stirred at room temperature for 1 hour. TLC showed the absence of starting material. The solution was concentrated to give the crude titled compound without further purification.

5

Step C: Preparation of Schiff Base:

To the mixture of step (9B) was added methanol (0.8 ml) and 20 mg. of *t*-butyl-*p*-aminobenzoate and 5 mg toluenesulfonic acid and the mixture stirred at room temperature for 18 hours, to yield the crude Schiff base.

10

Step D: Preparation of (\pm) cis-3-Ethyl-4-*t*-butoxycarbonylphenylaminomethyl-N-toluenesulfonyl-2-azetidinone

The product of step (9C) was hydrogenated at 40psi in the presence of 10% Pd/C. Flash chromatography gave the crude titled product.

15

Step E: Preparation of (\pm) cis-3-Ethyl-4-hydroxycarbonyl-phenylaminomethyl-N-toluenesulfonyl-2-azetidinone

The *tert*-butyl moiety was cleaved using 30% trifluoroacetic acid (TFA) in CH_2Cl_2 for 5 hours to give the titled compound.

20

Example 10

25

Preparation of (\pm)-3,3-Dimethyl-4-(1,2-epoxy-2-phenyl)ethyl-N-toluenesulfonyl-2-azetidinone

30

Step A: Preparation of (\pm)-3,3-Dimethyl-4-(1,2-epoxy-2-phenyl)ethyl-2-azetidinone

35

To 670 mg of (\pm)-3,3-Dimethyl-4-(β -styryl)-2-azetidinone in 10 ml of CH_2Cl_2 was added 616 mg of *m*-chloroperbenzoic acid. The mixture was stirred overnight, and filtered. The filtrate was purified via prep. tic on silica gel plates developed with CH_2Cl_2 :hexane (1:1). The impure product was further purified via prep. tic on silica gel plates developed with EtOAc:hexane (2:3) to give the titled compound. NMR (CDCl_3): δ 1.28 (3H, s), 1.40 (3H, s), 3.1 (1H, dd), 3.26 (1H, d), 3.69 (1H, d), 6.17 (1H, brs).

40

Step B Preparation of (\pm)-3,3-Dimethyl-4-(1,2-epoxy-2-phenyl)ethyl-N-toluenesulfonyl-2-azetidinone

45

To 25 mg (\pm)-3,3-Dimethyl-4-(1,2-epoxy-2-phenyl)ethyl-2-azetidinone in 0.3 ml of TBAB solution was added 120 mg of tosyl chloride and 20 mg of pulverized KOH. The mixture was stirred at room temperature overnight. The mixture was filtered and the filtrate purified via prep. tic on silica gel (1000 μ) developed with hexane: CH_2Cl_2 (1:4) to give the titled compound.

50

NMR (CDCl_3): δ 1.21 (3H, s), 1.25 (3H, s), 2.45 (3H, s), 3.06 (1H, dd), 3.51 (1H, d), 3.82 (1H, d), 7.2-7.5 (7H, m), 7.60 (2H, m).

Example 11

55

Preparation of (\pm)-3,3-Dimethyl-4-(1-benzoyloxy-2-phenyl)ethyl-N-toluenesulfonyl-2-azetidinone

Step A: Preparation of (±)-3,3-Dimethyl-4-(1-hydroxy-2-phenyl)ethyl-2-azetidinone

A solution of 64 mg of (±)-3,3-Dimethyl-4-(1,2-epoxy-2-phenyl)ethyl-2-azetidinone in 5 ml of EtOAc was hydrogenated at 1 atm. in the presence of palladium for 1 hour. The mixture was filtered through a celite pad washed with EtOAc/MeOH. The filtrate was concentrated to give the titled compound.
 NMR (CDCl₃) δ 1.28 (3H, s), 1.34 (3H, s), 2.64 (1H, dd), 2.86 (1H, dd), 3.30 (1H, d), 3.87 (1H, t), 6.05 (1H, s), 7.0-7.5 (5H, m).

Step B: Preparation of (±)-3,3-Dimethyl-4-(1-benzoyloxy-2-phenyl)ethyl-2-azetidinone


To 25 mg of the product of step (11A) in 0.5 ml of CH₂Cl₂ was added 0.2 ml of pyridine and then a solution of 0.1 ml of benzoyl chloride in 0.2 ml of CH₂Cl₂. The mixture was stirred at room temperature for 2 hours and concentrated. The residue was pumped to dryness and purified via prep. tlc on silica gel (1500μ) developed with 10% EtOAc in CH₂Cl₂ to give the titled compound.
 NMR (CDCl₃): δ 1.24 (3H, s), 1.33 (3H, s), 3.06 (2H, ddd), 3.58 (1H, d), 5.50 (1H, dd), 5.61 (1H, m), 7.1-7.7 (8H, m), 8.0 (2H, dd).

Step C: Preparation of (±)-3,3-Dimethyl-4-(1-benzoyloxy-2-phenyl)ethyl-N-toluenesulfonyl-2-azetidinone

To the product of step (11B) in 0.2 ml of TBAB solution was added 50 mg of tosyl chloride and 10 mg of pulverized KOH and the mixture stirred at room temperature overnight. The mixture was purified via prep. tlc on silica gel (1000μ) and developed with CH₂Cl₂ halfway and then 2% ETOAc in CH₂Cl₂ to give the titled compound.
 NMR (CDCl₃): δ 3.14 (3H, s), 3.15 (3H, s), 2.40 (3H, s), 3.24 (2H, ddd), 4.03 (1H, d), 5.36 (1H, m), 7.2-7.6 (10H, m), 7.70 (2H, dd), 7.89 (2H, d).

Claims

1. A compound of structural formula (I)



(I)


wherein:

Q is

- a. C₁₋₅ alkyl;
 - b. C₆₋₁₀ aryl or C₆₋₁₀ heteroaryl including one heteroatom selected from N, O, or S;
 - c. C₇₋₅ aralkyl or C₇₋₁₅ heteroaralkyl;
 - d. C₆₋₁₀ aryl or C₆₋₁₀ heteroaryl substituted with W;
 - e. C₇₋₁₅ aryl or C₇₋₁₅ heteroaryl wherein the aryl or heteroaryl moiety is substituted with W;
 - f. OH;
- R₃ and R₅ are independently selected from:
- a. H,
 - b. C₁₋₅ hydroxyalkyl,
 - c. C₁₋₅ alkyl,
 - d. C₁₋₅ alkoxy,
 - e. C₂₋₆ alkenyl;
- R₄ is CHR₀R;
- R₀ is H, C₁₋₅ alkyl, ph nyl or phenyl C₁₋₅ alkyl;


R is

- a. $O-(CO)_{p_2}-(A)_{p_1}-R_1$,
 - b. OSO_2R_1 ,
 - c. AR_1 ,
- 5 d. R together with R_0 and the C to which it is attached form



10

- e. R together with R_0 and the C to which it is attached form




15

p_1 and p_2 are independently 0 or 1; provided that p_1 and p_2 are not both 0;

20 R_1 is

- a. H,
- b. aryl, or a heteroaryl group of 5 to 10 atoms, including one to two heteroatoms, selected from:

25




30


35

- c. phenyl $S(O)_n$ wherein n is 0 to 2,
 - d. C_3-6 cycloalkyl;
- B is O, N or S;
A is

45



50



55

- c. NH;
- R_2 is
- a. H,

- b. C₁₋₅ alkoxy,
- c. C₁₋₅ alkyl,
- d. OH,
- e. OC(O)CH₃;
- 5 W is
 - a. C₁₋₅ alkyl,
 - b. C₁₋₅ alkoxy,
 - c. NO₂,
 - d. NHR₃,
 - 10 e. C₁₋₃ alkylS(O)p₃
 - f. halogen,
 - g. COOR₃,
 - h. OCOR₃,
 - i. NHCOR₃,
 - 15 j. NHSO₂R₃,
 - k. NHSO₂phenyl;
- X, Y, and Z are independently selected from:
 - a. H,
 - b. halogen,
 - 20 c. C₁₋₅ alkyl
 - d. C₁₋₅ alkoxy,
 - e. C₁₋₅ alkoxy carbonyl
 - f. C₁₋₅ alkyl carbonylamino,
 - g. amino,
 - 25 h. phenoxy,
 - i. phenyl C₁₋₅ alkoxy,
 - j. C₁₋₅ alkylSO₂NH,
 - k. NO₂,
 - l. diphenylmethyloxycarbonyl,
 - 30 m. C₁₋₅ alkylthio,
 - n. aminocarbonyl,
 - o. carboxy,
 - p. C₁₋₅ alkenyloxy
 - q. C₁₋₅ alkylamido,
 - 35 r. trifluoromethyl.
- 2. A compound of Claim 1 wherein: R is O-(CO)p₂-(A)p₁, R₁, and R₀ and R₅ are H.
- 3. A compound of Claim 2 wherein p₁ is 0 p₂ is 1 and Q is 4-methylphenyl, selected from the group where in:

40

45

50

55

and the
configuration at
the 3,4 position is

	R_2	R_4	
5			
	a. CH_3CH_2-	3-thienyl CO_2CH_2-	<i>cis</i>
	b. CH_3CH_2-	3-methanesulfonamidophenyl- CO_2CH_2-	<i>cis</i>
10	c. CH_3CH_2-	2-nitrophenyl CO_2CH_2-	<i>cis</i>
	d. CH_3CH_2-	4-pyridyl CO_2CH_2-	<i>cis</i>
	e. CH_3CH_2-	2-pyridyl CO_2CH_2-	<i>cis</i>
15	f. CH_3CH_2-	5-(2-methoxypyridyl) CO_2CH_2-	<i>cis</i>
	g. CH_3CH_2-	2-furanyl CO_2CH_2-	<i>cis</i>
	h. CH_3CH_2-	2-(5-n-butylpyridyl) CO_2CH_2-	<i>cis</i>
20	i. CH_3CH_2-	4-methoxyphenyl CO_2CH_2-	<i>cis</i>
	j. CH_3CH_2-	4-methoxyphenyl CO_2CH_2-	<i>cis</i>
	k. CH_3CH_2-	3-nitrophenyl CO_2CH_2-	3S, 4S
	l. CH_3CH_2-	2- $\text{CH}_3\text{SO}_2\text{NH}$ phenyl CO_2CH_2-	3R, 4R
25	m. CH_3CH_2-	4-pyridazinyl CO_2CH_2-	<i>cis</i>
	n. CH_3CH_2-	2- NH_2 phenyl CO_2CH_2-	<i>cis</i>
	o. CH_3CH_2-	3-furanyl CO_2CH_2-	<i>cis</i>
30	p. CH_3CH_2-	2-quinolinylnyl CO_2CH_2-	<i>cis</i>
	q. CH_3CH_2-	4-benzoxypyphenyl CO_2CH_2-	<i>cis</i>
	r. CH_3CH_2-	4-quinolinylnyl CO_2CH_2-	<i>cis</i>
35	s. CH_3CH_2-	2-thienyl CO_2CH_2-	<i>cis</i>
	t. CH_3CH_2-	3-(2-phenoxypyridyl) CO_2CH_2-	<i>cis</i>
40			
45			
50			
55			

and the
configuration at
the 3,4 position is

	R_3	R_4	
5			
	u. CH_3CH_2-	3- $\text{CH}_3\text{CONHphenylCO}_2\text{CH}_2-$	<u>cis</u>
	v. CH_3CH_2-	3- $\text{NH}_2\text{phenylCO}_2\text{CH}_2-$	<u>cis</u>
10	w. CH_3CH_2-	4-((Ph) $_2\text{COOC}$)phenylCO $_2\text{CH}_2-$	<u>cis</u>
	x. CH_3CH_2-	4- $\text{HOOCphenylCO}_2\text{CH}_2-$	<u>cis</u>
	y. CH_3CH_2-	4- $\text{NO}_2\text{phenylCO}_2\text{CH}_2-$	<u>cis</u>
15	z. $\text{CH}_3\text{CH}_2\text{CH}_2-$	4- $\text{CH}_3\text{OphenylCO}_2\text{CH}_2-$	<u>cis</u>
	aa. HOCH_2-	4- $\text{CH}_3\text{OphenylCO}_2\text{CH}_2-$	<u>cis</u>
	bb. CH_3CH_2-	3-pyridylCO $_2\text{CH}_2-$	<u>cis</u>
	cc. CH_3CH_2-	4- $\text{CH}_3\text{OphenylCO}_2\text{CH}_2-$	<u>cis</u>
20	dd. CH_3CH_2-	phenylCO $_2\text{CH}_2-$	<u>cis</u>
	ee. CH_3CH_2-	3-CoumarylCO $_2\text{CH}_2-$	<u>cis</u>
	ff. CH_3CH_2-	2-benzofuranylCO $_2\text{CH}_2-$	<u>cis</u>
25	gg. CH_3CH_2-	2-furanylCO $_2\text{CH}_2-$	S,S
	hh. $\text{CH}_3\text{CHCH}_3-$	4-methoxyphenylCO $_2\text{CH}_2-$	<u>cis</u>
	ii. CH_3CH_2-	4- $\text{CH}_3\text{O}_2\text{CphenylCO}_2\text{CH}_2-$	<u>cis</u>
30	jj. CH_3CH_2-	4-n-butylcarbamoylphenylCO $_2\text{CH}_2-$	<u>cis</u>
	kk. CH_3CH_2-	4-aminophenylCO $_2\text{CH}_2-$	<u>cis</u>
	ll. CH_3CH_2-	3-methoxy-4-allyloxy-5-nitro- phenylCO $_2\text{CH}_2-$	<u>cis</u>
35	mm. CH_3CH_2-	3-methoxy-4-allyloxy-5-(methyl- amido)phenylCO $_2\text{CH}_2-$	<u>cis</u>
	nn. CH_3CH_2-	4-methanesulfonamidophenylCO $_2\text{CH}_2-$	<u>cis</u>
40	oo. CH_3CH_2-	5-(2,3-dimethoxypyridyl)CO $_2\text{CH}_2-$	<u>cis</u>
	pp. CH_3CH_2-	6-(2,3-dimethoxypyridyl)CO $_2\text{CH}_2-$	<u>cis</u>
	qq. CH_3CH_2-	4-(2-methoxythiazolyl)CO $_2\text{CH}_2-$	<u>cis</u>
45	rr. $\text{CH}_3\text{CH}_2\text{CH}_2-$	4- $\text{CH}_3\text{OphenylCO}_2\text{CH}_2-$	<u>trans</u>
	ss. HOCH_2-	4- $\text{CH}_3\text{OphenylCO}_2\text{CH}_2-$	<u>trans</u>
	tt. CH_3CH_2-	4- $\text{CH}_3\text{OphenylCO}_2\text{CH}_2-$	<u>trans</u>

50 4. A compound of Claim 2 wherein p_1 is 1 and p_2 is 1, selected from the group where in:


55

	<u>R₃</u>	<u>R₄</u>	<u>and the configuration at the 3,4 position is</u>
5	a. CH ₃ CH ₂ -	4-CH ₃ OphenylC = CCO ₂ CH ₂ -	<u>cis</u>
	b. CH ₃ CH ₂ -	4-CH ₃ OphenylCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	c. CH ₃ CH ₂ -	3-pyridylCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	d. CH ₃ CH ₂ -	phenyl-(R)-CHOCH ₃ CO ₂ CH ₂ -	<u>S,S</u>
	e. H	phenyl-(S)-CHOCH ₃ CO ₂ CH ₂ -	<u>4S</u>
	f. H	phenyl-(R)-CHOCH ₃ CO ₂ CH ₂ -	<u>4S</u>
10	g. CH ₃ CH ₂ -	phenyl-(R)-CHOCH ₃ CO ₂ CH ₂ -	<u>cis</u>
	h. CH ₃ CH ₂ -	3-pyridylC = CCO ₂ CH ₂ -	<u>cis</u>
	i. CH ₃ CH ₂ -	3-pyridylCH ₂ CO ₂ CH ₂ -	<u>3S,4S</u>
	j. CH ₃ CH ₂ -	4-methoxyphenylCH ₂ CO ₂ CH ₂ -	<u>3S,4S</u>
	k. CH ₃ CH ₂ -	4-methoxyphenylC = CCO ₂ CH ₂ -	<u>3S,4S</u>
15	l. CH ₃ CH ₂ -	2-furanylCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	m. CH ₃ CH ₂ -	2-furanylC = CCO ₂ CH ₂ -	<u>cis</u>
	n. CH ₃ CH ₂ -	phenoxyCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	o. CH ₃ CH ₂ -	3-methyl-4-ethoxyphenylCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	p. CH ₃ CH ₂ -	2-pyridylCH ₂ CO ₂ CH ₂ -	<u>cis</u>
20	q. CH ₃ CH ₂ -	phenylthioCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	r. CH ₃ CH ₂ -	4-ethoxyphenylCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	s. CH ₃ CH ₂ -	3,4,5-trimethoxyphenylCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	t. CH ₃ CH ₂ -	3,4,5-trimethoxyphenylC = CCO ₂ CH ₂ -	<u>cis</u>
	u. CH ₃ CH ₂ -	3-bromo-4-propanoxy-5-methoxyphenylC = CCO ₂ CH ₂ -	<u>cis</u>
25	v. CH ₃ CH ₂ -	4-methylphenylaminoCO ₂ CH ₂ -	<u>cis</u>
	w. CH ₃ CH ₂ -	cyclohexylaminoCO ₂ CH ₂ -	<u>cis</u>
	x. CH ₃ CH ₂ -	4-trifluoromethylphenylamino-CO ₂ CH ₂ -	<u>cis</u>
	y. CH ₃ CH ₂ -	3,4-dimethoxyphenylCH ₂ CO ₂ CH ₂ -	<u>cis</u>
	z. CH ₃ CH ₂ -	6-(2,3-dimethoxypyridyl)C = CCO ₂ CH ₂ -	<u>cis</u>
30	aa. CH ₃ CH ₂ -	4-fluorophenylaminoCO ₂ CH ₂ -	<u>cis</u>

5. A compound of Claim 1 wherein: R is OSO₂R₁ and R₀ and R₅ are H, selected from the group wherein:

<u>R₃ is</u>	<u>R₄ is</u>	<u>and the configuration at the 3,4 position is</u>
a. CH ₃ CH ₂ -	4-CH ₃ phenylISO ₂ OCH ₂ -	<u>cis</u>

6. A compound of Claim 1 wherein: R is O-(CO)_{p2}-(A)_{p1}-R₁, AR₁ or R together with R₀ and the C to which it is attached form




P₁ and P₂ are independently 0 or 1, provided that P₁ and P₂ are not both zero;

R₀ is H, CH₃ or phenyl; provided that when R is O-(CO)_{p2}-(A)_{p1}-R₁, R₀ is CH₃ or phenyl;

R₅ is H.

7. A compound of Claim 6 wherein:



R_0 is H, or CH_3 , selected from the group wherein:

	<u>R_3 is</u>	<u>R_4 is</u>	<u>and the configuration at the 3,4 position is</u>
a.	$\text{CH}_3\text{CH}_2\text{CH}_2-$	phenylC = C-	trans
b.	$\text{CH}_3\text{CH}_2\text{CH}_2-$	phenylC = C-	cis
c.	CH_3CH_2-	phenylC = C-	trans
d.	HOCH_2-	phenylC = C-	cis
e.	CH_3CH_2-	phenylC = CCH ₃	trans
f.	CH_3CH_2-	phenylC = CCH ₃	cis

8. A compound of Claim 6 wherein:

R is $\text{O}-(\text{CO})_{\text{p}_2}-(\text{A})_{\text{p}_1}-\text{R}_1$ or AR_1 ;

R_0 is H, CH_3 or phenyl; provided that when R is $\text{O}-(\text{CO})_{\text{p}_2}-(\text{A})_{\text{p}_1}-\text{R}_1$, R_0 is CH_3 or phenyl.

9. A compound of Claim 8 wherein:

P_1 is 0, selected from the group wherein:

and the
configuration at
the 3,4 position is

<u>R_3 is</u>	<u>R_4 is</u>	
-----------------------------------	-----------------------------------	--


a.	CH_3CH_2-	4-HO ₂ CphenylNHCH ₂ -	cis
b.	CH_3CH_2-	phenylCH ₂ CH ₂ -	cis
c.	CH_3CH_2-	4-CH ₃ OphenylCO ₂ CHCH ₃	cis
d.	CH_3CH_2-	phenylCH ₂ CHCH ₃	cis
e.	CH_3CH_2-	4-CH ₃ OphenylCO ₂ CHphenyl	cis
f.	H	phenylCHCHCH ₃ OCOCH ₃	
g.	H	CH ₃ CO ₂ CH ₂ CHCH ₃	

10. A compound of Claim 1 wherein:

R_0 is H, CH_3 or CH_2phenyl ;

R_5 is C_{1-5} alkyl or C_{1-5} alkoxy;

R is $\text{O}-(\text{CO})_{\text{p}_2}-(\text{A})_{\text{p}_1}-\text{R}_1$, AR_1 or R together with R_0 and the C to which it is attached form




P_1 and P_2 are independently 0 or 1;

provided that P_1 and P_2 are not both zero; selected from the group wherein:

	<u>R₃ is</u>	<u>R₅ is</u>	<u>R₄ is</u>
5	a. CH ₃	CH ₃	phenylCH ₂ CH ₂ -
	b. CH ₃	CH ₃	phenylCO ₂ CHCH ₂ phenyl
	c. CH ₃	CH ₃	phenylCH=CH-
10	d. CH ₃	CH ₃	phenylCH-CH-
			O

11. A hypocholesterolemic, hypolipidemic pharmaceutical composition comprising a pharmaceutically acceptable carrier and a therapeutically effective amount of a compound as defined in Claim 1.



Europäisches Patentamt
European Patent Office
Office européen des brevets



Publication number:

0 415 487 A3

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 90202269.8

(51) Int. Cl.⁵: C07D 205/08, C07D 401/12,
C07D 405/06, A61K 31/395

(22) Date of filing: 24.08.90

(30) Priority: 31.08.89 US 401391

(43) Date of publication of application:
06.03.91 Bulletin 91/10

(94) Designated Contracting States:
CH DE FR GB IT LI NL

(38) Date of deferred publication of the search report:
11.03.92 Bulletin 92/11


(71) Applicant: MERCK & CO. INC.
126, East Lincoln Avenue P.O. Box 2000
Rahway New Jersey 07065-0900(US)

(72) Inventor: Yang, Shu Shu
324 Eileen Way
Bridgewater, NJ 08807(US)
Inventor: Chiang, Yuan-Ching P.
73 Mitchell Avenue
Piscataway, NJ 08854(US)
Inventor: Heck, James V.
961 Nepawin Lane
Scotch Plains, NJ 07076(US)
Inventor: Chang, Michael N.
2970 Windy Bush Road
Newton, PA 18940(US)

(74) Representative: Hesketh, Alan, Dr. et al
European Patent Department Merck & Co.,
Inc. Terlings Park Eastwick Road
Harlow Essex, CM20 2QR(GB)

(54) Beta-lactams as anticholesterolemic agents.

(57) Compounds of formula (I), which are useful as
antihypercholesterolemic agents, are disclosed.



(I)

EP 0 415 487 A3

THIS PAGE BLANK (USPTO)



European
Patent Office

EUROPEAN SEARCH REPORT

Application Number

EP 90 20 2269

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.5)
A	EP-A-0 293 132 (MERCK & CO., INC.) "claims"	1,11	C 07 D 205/08 C 07 D 401/12 C 07 D 405/06

D,A	EP-A-0 293 128 (MERCK & CO., INC.) "claims"	1,11	A 61 K 31/395

D,A	EP-A-0 285 254 (MERCK & CO., INC.) "claims"	1,11	

A	EP-A-0 199 630 (MERCK & CO., INC.) "claims"	1,11	

The present search report has been drawn up for all claims			
Place of search		Date of completion of search	Examiner
The Hague		07 January 92	CHOULY J.
CATEGORY OF CITED DOCUMENTS			
X: particularly relevant if taken alone		E: earlier patent document, but published on, or after the filing date	
Y: particularly relevant if combined with another document of the same category		D: document cited in the application	
A: technological background		L: document cited for other reasons	
O: non-written disclosure		&: member of the same patent family, corresponding document	
P: intermediate document			
T: theory or principle underlying the invention			

THIS PAGE BLANK (USPTO)
